



WELL-POSEDNESS CHARACTERIZATIONS FOR SYSTEM OF MIXED HEMIVARIATIONAL INEQUALITIES

KARTIKESWAR MAHALIK AND CHANDAL NAHAK

Received 11 April, 2022

Abstract. In this paper, we concern with the concept of well-posedness and well-posedness in generalized sense for a system of mixed hemivariational inequality (SMHVI) with perturbations. We establish several metric characterizations and equivalent conditions of well-posedness for SMHVI. Our main results improve and extend some announced work in the literature.

2010 Mathematics Subject Classification: 49J40; 47J20

Keywords: system of mixed hemivariational inequalities, approximating sequence, well-posedness, perturbation

1. INTRODUCTION

The theory of well-posedness in optimization problems and various comprehensive problems correlated to it such as variational inequalities, fixed point problems, equilibrium problems, saddle point problems, and inclusion problems has been paid deep impact attention by researchers due to their essential contributions to physics, mechanics, engineering, and economics. Tykhonov [20] first proposed the concept of well-posedness for the unconstrained global minimization problems, i.e., a minimization problem is well-posed, if it possesses a unique minimizer and the limit of every subsequence of the minimizing sequence is the unique minimizer. Indeed, an optimization problem may not have a unique solution. Hence, the notion of generalized well-posedness was introduced, which enhances the precise solution of the problem, under some mild conditions, by ensuring the convergence of every approximating solutions to the exact solution. Accordingly, the approximating approach captured the concept of well-posedness that plays a vital aspect in well-posedness theory. For more related research works concerning to the well-posedness results

The first author was supported by the Council of Scientific and Industrial Research (CSIR), India, under grant No. 09/081(1306)/2017-EMR-I. The second author was partially supported by the SERB, DST, Govt. of India, Grant No MTR/2022/000296.

© 2024 The Author(s). Published by Miskolc University Press. This is an open access article under the license [CC BY 4.0](#).

of optimization problems, we refer to [11, 14]. Later, the notion of well-posedness turned toward variational inequalities in different aspects. Lucchetti and Patrone [17] first studied and developed the concept of well-posedness for variational inequalities by Ekeland's technique. From then on, researchers are dedicated to investigate the well-posedness for variational inequalities and build their links with the well-posedness of inclusion and minimization problems. Lignola [13] and Lignola and Morgan [14] coined the notion of L -well-posedness for quasivariational inequalities and parametric well-posedness for variational inequalities, respectively. Ceng et al. [3] explored various necessary and sufficient conditions of the well-posedness for the mixed quasivariational-like inequalities. Meanwhile, Fang et al. [8, 9] demonstrated different forms of well-posedness results for mixed variational inequalities, which were further generalized by Ceng et al. [2]. In the works of [4, 8, 9], the authors discussed various necessary and sufficient metric characterizations of well-posedness. There, the authors also elaborated on the corresponding inclusion problems as well as the corresponding fixed point problems and established the equivalent relations between the well-posedness of mixed variational inequalities and their corresponding inclusion problems.

In the last decade, the study of well-posedness devoted special attention to hemivariational inequality problem (HVIP), which is a powerful and important generalization of variational inequalities. The HVIP was first investigated and explored by Panagiotopoulos [19] at the beginning of 1980s to formulate several classes of unilateral mechanical problems with nonsmooth and nonconvex energy functional. The HVIP is described through Clarke's generalized gradient for nonconvex and nondifferentiable functions. It is straightforward to see that if the involved functions are convex, then the HVIP weakens to the variational inequality (see [6]). Consequently, several extensive well-posedness results (for e.g. see, [2, 5, 10, 15, 22–24]) have been investigated and widely discussed for HVIP. Goeleven and Motreanu [10] first considered the well-posedness consequences for HVIP. Xiao et al. [23] discussed the well-posedness characterizations for a class of HVIP. Recently, Xiao et al. [24] also explored the well-posedness results for hemivariational inequalities by utilizing the notion of strongly relaxed monotonicity. By introducing $\alpha - \eta$ monotone mappings, Liu et al. [15] established $L - \alpha$ well-posedness for mixed quasivariational hemivariational inequality through Mosco convergence of sets. Xiao and Huang [22] proposed the well-posedness results for variational-hemivariational inequalities with perturbations. In 2013, Costea and Varga [7] considered the existence results for the system of hemivariational inequality and also derived the Nash generalized derivative points. Recently, Wang et al. [21] generalized the concept of well-posedness to a system of hemivariational inequalities in product Banach space. Besides, the authors in [21–23] also built the equivalence results between the well-posedness of the inequality problems and the well-posedness of its derived inclusion problems. Further, Ceng et al. [5]

studied the well-posedness results for system of time-dependent hemivariational inequalities. However, the research works based on the well-posedness of solutions for the system of hemivariational inequalities are quite limited due to the complexity of its structure. The aim of this work is to develop the theoretical framework on well-posedness for a new class of system of mixed hemivariational inequality with perturbations.

Motivated by the aforesaid research works on well-posedness, in this work, our aim is to introduce and generalize the concept of well-posedness for the SMHVI with perturbations, which includes the HVIP as a special case. Section 2 briefly revisits some preliminary definitions and notations, which will be used to achieve the main results. Section 3 formulates the SMHVI with perturbations and the concept of approximating sequence as well as strong (resp., weak) well-posedness. Section 4 explores the necessary theorems and the metric characterizations for the well-posedness of SMHVI.

2. PRELIMINARIES

Let X be a Banach space with dual space X^* . We signify $\langle \cdot, \cdot \rangle_{X^* \times X}$ as the duality pairing between X^* and X . We denote \rightarrow , \rightharpoonup and $\xrightarrow{w^*}$ as the strong, weak and weak* convergence, respectively, in the appointed space.

Definition 1 ([18, Definition 1.31]). A sequence $\{x_n\} \subset X$ is weakly convergent iff there exists $x \in X$ s.t.

$$\langle x^*, x_n \rangle_{X^* \times X} \rightarrow \langle x^*, x \rangle_{X^* \times X}, \quad \forall x^* \in X^*.$$

Definition 2 ([18, Definition 1.42]). A sequence of functional $\{x_n^*\} \subset X^*$ is called weakly* convergent to $x^* \in X^*$ iff

$$\langle x_n^*, x \rangle_{X^* \times X} \rightarrow \langle x^*, x \rangle_{X^* \times X}, \quad \forall x \in X.$$

Theorem 1 ([18, Proposition 1.37 and Proposition 1.45]). *If $\{x_n\} \subset X$ and $\{x_n^*\} \subset X^*$ with $x_n \rightarrow x$ (resp. $x_n \rightharpoonup x$) in X and $x_n^* \xrightarrow{w^*} x^*$ (resp. $x_n^* \rightarrow x^*$) in X^* , then*

$$\langle x_n^*, x_n \rangle_{X^* \times X} \rightarrow \langle x^*, x \rangle_{X^* \times X}.$$

A function $\phi: X \rightarrow \mathbb{R} \cup \{\infty\}$ is called lower semicontinuous (l.s.c.) if it fulfills

$$\phi(x) \leq \liminf_{n \rightarrow \infty} \phi(x_n)$$

for all sequence $x_n \in X$ with $x_n \rightarrow x$.

Definition 3 ([21, Definition 2.13]). We say that a functional $J: X \rightarrow \mathbb{R}$ is locally Lipschitz, if for every $z \in X$, there exists a neighborhood V of z and a constant $K_z > 0$ s.t. $|J(x) - J(y)| \leq K_z \|x - y\|_X$ for all $x, y \in V$.

Definition 4 ([21, Definition 2.16]). Let $J: X \rightarrow \mathbb{R}$ be a locally Lipschitz functional. We denote $J^\circ(x; y)$ and $\bar{\partial}J(x)$ as the Clarke's generalized directional derivative at the point $x \in X$ in the direction $y \in X$ and Clarke's generalized gradient of J at $x \in X$, respectively, which are defined as:

$$J^\circ(x; y) = \limsup_{w \rightarrow x, \varepsilon \rightarrow 0^+} \frac{J(w + \varepsilon y) - J(w)}{\varepsilon},$$

and $\bar{\partial}J(x) = \{\xi \in X^* : \langle \xi, y \rangle_{X^* \times X} \leq J^\circ(x; y), \forall y \in X\}$, respectively.

Lemma 1 ([6, Proposition 2.1.1 and Proposition 2.1.2]). *Let $J: X \rightarrow \mathbb{R}$ be a locally Lipschitz functional of rank L . Then*

- (i) *the function $y \rightarrow J^\circ(x; y)$ is finite, homogenous, subadditive on X and satisfies $|J^\circ(x; y)| \leq L\|y\|_X$;*
- (ii) *the functional $(x, y) \rightarrow J^\circ(x, y)$ is upper semicontinuous;*
- (iii) *$J^\circ(x; -y) = (-J)^\circ(x; y)$;*
- (iv) *for each $x \in X$, $\bar{\partial}J(x)$ is a nonempty, convex, weak*-compact subset of X^* and $\|\xi\|_{X^*} \leq L$ for all $\xi \in \bar{\partial}J(x)$;*
- (v) *for each $y \in X$, one has $J^\circ(x; y) = \max_{\xi \in \bar{\partial}J(x)} \langle \xi, y \rangle_{X^* \times X}$.*

Definition 5 ([22, Definition 2.3]). The measure of non-compactness for the set $D \subset X$ is defined by

$$\mu(D) = \inf \left\{ \gamma > 0 : D \subset \bigcup_{i=1}^n D_i, \text{diam}|D_i| < \gamma, i = 1, 2, \dots, n \right\},$$

where $\text{diam}|D_i|$ denotes the diameter of the set D_i .

If D is a bounded subset of X then $\mu(D) = 0$ iff \bar{D} is compact. Also, if D_1 is a subset of D_2 then $\mu(D_1) \leq \mu(D_2)$.

Definition 6 ([16, Definition 8.1.7]). Let C and D be two nonempty subsets of X . The Hausdorff distance $\mathcal{H}(\cdot, \cdot)$ between C and D is defined by

$$\mathcal{H}(C, D) = \max\{e(C, D), e(D, C)\},$$

where $e(C, D) = \sup_{c \in C} d(c, D)$ with $d(c, D) = \inf_{d \in D} \|c - d\|_X$.

Here, we want to emphasize that if $\{C_n\}$ be a sequence of nonempty subsets of X . Then we say that C_n converges to C with respect to $\mathcal{H}(\cdot, \cdot)$ iff $\mathcal{H}(C_n, C) \rightarrow 0$.

Definition 7 ([22, Definition 2.1 and Definition 2.2]). Let $G: X \rightarrow X^*$ be a single-valued operator, then G is

- (i) *hemicontinuous, iff for all $x, y \in X$, the function $\varepsilon \rightarrow \langle G(x + \varepsilon(y - x)), y - x \rangle_{X^* \times X}$ from $[0, 1]$ into \mathbb{R} is continuous at 0^+ ;*
- (ii) *monotone, iff $\langle G(x) - G(y), x - y \rangle_{X^* \times X} \geq 0, \forall x, y \in X$.*

- (H_j) $J_i: X_i \rightarrow \mathbb{R}$ is locally Lipschitz functional,
 (H_ϕ) $\phi_i: X_i \rightarrow \mathbb{R} \cup \{\infty\}$ is proper, convex and l.s.c. map with effective domain:
 $\text{dom}(\phi_i) = \{x_i \in X_i : \phi_i(x_i) < \infty\} \neq \emptyset$,
 (H_c) $S_i: X_1 \times \cdots \times X_i \times \cdots \times X_k \rightarrow X_i^*$ is continuous on $X_1 \times \cdots \times X_i \times \cdots \times X_k$,
 (H_h) $A_i: X_1 \times \cdots \times X_i \times \cdots \times X_k \rightarrow X_i^*$ is hemicontinuous with respect to X_i .

We end this section by addressing the following equivalence results between the sets $\Upsilon(\rho)$ and $\Psi(\rho)$ for any $\rho > 0$.

Lemma 2. *For each $i \in \{1, \dots, k\}$, suppose that $A_i: X_1 \times \cdots \times X_i \times \cdots \times X_k \rightarrow X_i^*$ satisfies the hypothesis (H_m) and (H_h). Further assume that for each $i \in \{1, \dots, k\}$, $\phi_i: X_i \rightarrow \mathbb{R} \cup \{\infty\}$ is a proper and convex functional, and $J_i: X_i \rightarrow \mathbb{R}$ satisfies the hypothesis (H_j). Then $\Upsilon(\rho) = \Psi(\rho)$ for any $\rho > 0$.*

Proof. Let $x = (x_1, \dots, x_i, \dots, x_k) \in X_1 \times \cdots \times X_i \times \cdots \times X_k$ and $x \in \Upsilon(\rho)$. Thus, for each $i \in \{1, \dots, k\}$, we have

$$\begin{aligned} & \langle A_i(x_1, \dots, x_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) \\ & + \phi_i(y_i) - \phi_i(x_i) \geq -\rho \|y_i - x_i\|_{X_i}, \quad \forall y_i \in X_i. \end{aligned} \quad (3.3)$$

Since A_i is monotone with respect to X_i (by H_m). We have

$$\langle A_i(x_1, \dots, y_i, \dots, x_k) - A_i(x_1, \dots, x_i, \dots, x_k), y_i - x_i \rangle_{X_i^* \times X_i} \geq 0, \quad \forall y_i, x_i \in X_i,$$

i.e.,

$$\begin{aligned} & \langle A_i(x_1, \dots, y_i, \dots, x_k), y_i - x_i \rangle_{X_i^* \times X_i} \\ & \geq \langle A_i(x_1, \dots, x_i, \dots, x_k), y_i - x_i \rangle_{X_i^* \times X_i}, \quad \forall x_i, y_i \in X_i. \end{aligned} \quad (3.4)$$

By combining (3.3) and (3.4), we get

$$\begin{aligned} & \langle A_i(x_1, \dots, y_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) \\ & + \phi_i(y_i) - \phi_i(x_i) \geq -\rho \|y_i - x_i\|_{X_i}, \quad \forall y_i \in X_i. \end{aligned}$$

i.e., $x = (x_1, \dots, x_i, \dots, x_k) \in \Psi(\rho)$, so, $\Upsilon(\rho) \subset \Psi(\rho)$.

Conversely, let $x = (x_1, \dots, x_i, \dots, x_k) \in X_1 \times \cdots \times X_i \times \cdots \times X_k$ and $x \in \Psi(\rho)$ for any $\rho > 0$. So, we yield that for each $i \in \{1, \dots, k\}$

$$\begin{aligned} & \langle A_i(x_1, \dots, y_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) \\ & + \phi_i(y_i) - \phi_i(x_i) \geq -\rho \|y_i - x_i\|_{X_i}, \quad \forall y_i \in X_i. \end{aligned} \quad (3.5)$$

Now for any $w = (w_1, \dots, w_i, \dots, w_k) \in X_1 \times \cdots \times X_i \times \cdots \times X_k$ and $t \in (0, 1)$, let

$$z_i = x_i + t(w_i - x_i) = tw_i + (1-t)x_i \in X_i, \quad i \in \{1, \dots, k\}.$$

Inserting z_i in (3.5), we have that for each $i \in \{1, \dots, k\}$

$$\begin{aligned} & \langle A_i(x_1, \dots, x_i + t(w_i - x_i), \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, t(w_i - x_i) \rangle_{X_i^* \times X_i} \\ & + J_i^\circ(x_i; t(w_i - x_i)) + \phi_i(x_i + t(w_i - x_i)) - \phi_i(x_i) \geq -\rho t \|w_i - x_i\|_{X_i}. \end{aligned}$$

By the convexity of ϕ_i and Lemma 1, we estimate that for each $i \in \{1, \dots, k\}$

$$\begin{aligned} & \langle A_i(x_1, \dots, x_i + t(w_i - x_i), \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, w_i - x_i \rangle_{X_i^* \times X_i} \\ & + J_i^\circ(x_i; w_i - x_i) + \phi_i(w_i) - \phi_i(x_i) \geq -\rho \|w_i - x_i\|_{X_i}. \end{aligned} \quad (3.6)$$

From (H_h) , A_i is hemicontinuous with respect to X_i for each $i \in \{1, \dots, k\}$. Taking the limit as $t \rightarrow 0$ in (3.6), we achieve

$$\begin{aligned} & \langle A_i(x_1, \dots, x_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, w_i - x_i \rangle_{X_i^* \times X_i} \\ & + J_i^\circ(x_i; w_i - x_i) + \phi_i(w_i) - \phi_i(x_i) \geq -\rho \|w_i - x_i\|_{X_i}, \quad \forall w_i \in X_i. \end{aligned}$$

Hence, the arbitrariness of w implies that $x \in \Upsilon(\rho)$. Thus, $\Psi(\rho) \subset \Upsilon(\rho) \forall \rho > 0$. \square

Remark 2. Lemma 2 improves and generalizes Lemma 3.1 in [22] and Lemma 3.1 in [23].

4. MAIN RESULTS

This section presents the metric characterizations and deduces some criteria, under which, SMHVI is strongly well-posed.

Proposition 1. For each $i \in \{1, \dots, k\}$, let X_i be a real Banach space with dual space X_i^* . Further, assume that for each $i \in \{1, \dots, k\}$, $A_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$, $S_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$, $J_i: X_i \rightarrow \mathbb{R}$ and $\phi_i: X_i \rightarrow \mathbb{R} \cup \{\infty\}$ satisfy the hypothesis (H_a) , (H_c) , (H_j) and (H_ϕ) , respectively. Then, $\Psi(\rho)$ is closed in $X_1 \times \dots \times X_i \times \dots \times X_k$ for any $\rho > 0$.

Proof. Let $x^n = (x_1^n, \dots, x_i^n, \dots, x_k^n) \in \Psi(\rho)$ be a sequence s.t. $(x_1^n, \dots, x_i^n, \dots, x_k^n) \rightarrow x = (x_1, \dots, x_i, \dots, x_k)$. Then for each $i \in \{1, \dots, k\}$,

$$\begin{aligned} & \langle A_i(x_1^n, \dots, y_i, \dots, x_k^n) + S_i(x_1^n, \dots, x_i^n, \dots, x_k^n) - f_i, y_i - x_i^n \rangle_{X_i^* \times X_i} + J_i^\circ(x_i^n; y_i - x_i^n) \\ & + \phi_i(y_i) - \phi_i(x_i^n) \geq -\rho \|y_i - x_i^n\|_{X_i}, \quad \forall y_i \in X_i. \end{aligned} \quad (4.1)$$

Since A_i, S_i, J_i , and ϕ_i , satisfy $(H_a), (H_c), (H_j)$ and (H_ϕ) , respectively. Thus, for each $i \in \{1, \dots, k\}$, by using (H_a) and Theorem 1, we obtain that

$$\langle A_i(x_1^n, \dots, y_i, \dots, x_k^n), y_i - x_i^n \rangle_{X_i^* \times X_i} \rightarrow \langle A_i(x_1, \dots, y_i, \dots, x_k), y_i - x_i \rangle_{X_i^* \times X_i}. \quad (4.2)$$

Also, taking the upper limit as $n \rightarrow \infty$ in (4.1), it follows from $(H_c), (H_j), (H_\phi)$ and (4.2) that for each $i \in \{1, 2, \dots, k\}$

$$\begin{aligned} & \langle A_i(x_1, \dots, y_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) \\ & + \phi_i(y_i) - \phi_i(x_i) \geq -\rho \|y_i - x_i\|_{X_i}, \quad \forall y_i \in X_i, \end{aligned}$$

which confirms that $x = (x_1, \dots, x_i, \dots, x_k) \in \Psi(\rho)$. Therefore, $\Psi(\rho)$ is closed. \square

The following result can be obtained by applying Lemma 2 and Proposition 1.

Corollary 1. For each $i \in \{1, \dots, k\}$, let X_i be real Banach space with dual space X_i^* and assume that $A_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$ satisfies the hypothesis (H_a) , (H_h) and (H_m) . Further, suppose that for each $i \in \{1, \dots, k\}$, $S_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$, $J_i: X_i \rightarrow \mathbb{R}$ and $\phi_i: X_i \rightarrow \mathbb{R} \cup \{\infty\}$ satisfy the hypothesis (H_c) , (H_j) and (H_ϕ) , respectively. Then $\Upsilon(\rho)$ is closed for any $\rho > 0$.

We now present the main results concerning to the strongly well-posedness of SMHVI.

Theorem 3. For each $i \in \{1, \dots, k\}$, let X_i be a real Banach space with dual space X_i^* and assume that $A_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$ satisfies the hypothesis (H_a) , (H_m) and (H_h) . Further, suppose that for each $i \in \{1, \dots, k\}$, $S_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$, $J_i: X_i \rightarrow \mathbb{R}$ and $\phi_i: X_i \rightarrow \mathbb{R} \cup \{\infty\}$ satisfy the hypothesis (H_c) , (H_j) and (H_ϕ) , respectively. Then SMHVI is strongly well-posedness iff

$$\Upsilon(\rho) \neq \emptyset \quad \forall \rho > 0 \quad \text{and} \quad \text{diam}(\Upsilon(\rho)) \rightarrow 0 \quad \text{as} \quad \rho \rightarrow 0.$$

Proof. Let SMHVI be strongly well-posed. Then SMHVI has a unique solution, say, $x = (x_1, \dots, x_i, \dots, x_k)$ and hence $x \in \Upsilon(\rho) \quad \forall \rho > 0$, which relates to $Q \neq \emptyset$ and $\Upsilon(\rho) \neq \emptyset \quad \forall \rho > 0$. Also, suppose that $\text{diam}(\Upsilon(\rho)) \not\rightarrow 0$ as $\rho \rightarrow 0$. Thus, there exists $\eta > 0$ and a sequence $\{\rho_n\} \subset \mathbb{R}_+$ with $\rho_n \rightarrow 0$ as $n \rightarrow \infty$ s.t. for each $n \in \mathbb{N}$, there exists $w^n = (w_1^n, \dots, w_i^n, \dots, w_k^n)$ and $z^n = (z_1^n, \dots, z_i^n, \dots, z_k^n)$ with $w^n, z^n \in \Upsilon(\rho_n)$ s.t.

$$\|w^n - z^n\| > \eta. \tag{4.3}$$

On the other hand, since $w^n, z^n \in \Upsilon(\rho_n)$, we have that the sequences $\{w^n\}$ and $\{z^n\}$ are both approximating sequences for SMHVI. Further, SMHVI is strongly well-posed, so the sequence $\{w^n\}$ and $\{z^n\}$ converges strongly to $x \in Q$, which contradicts (4.3).

In order to prove the sufficiency, let $x^n = (x_1^n, \dots, x_i^n, \dots, x_k^n)$ be an approximating sequence for SMHVI. Thus, there exists a sequence $\{\rho_n\} \subset \mathbb{R}_+$ with $\rho_n \rightarrow 0$ s.t. (3.1) holds, i.e., $x^n \in \Upsilon(\rho_n)$. Since $\text{diam}(\Upsilon(\rho_n)) \rightarrow 0$ as $\rho_n \rightarrow 0$, it is easy to draw that $\{x^n\}$ is a Cauchy sequence in $X_1 \times \dots \times X_i \times \dots \times X_k$. Since $X_1 \times \dots \times X_i \times \dots \times X_k$ is complete, and so $\{x^n\}$ converges strongly to some point $x = (x_1, \dots, x_i, \dots, x_k) \in X_1 \times \dots \times X_i \times \dots \times X_k$. Using the hypothesis (H_a) , (H_j) , (H_c) , (H_ϕ) and Lemma 1, it follows that for each $i \in \{1, \dots, k\}$,

$$\begin{aligned} & \langle A_i(x_1, \dots, y_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) \\ & + \phi_i(y_i) - \phi_i(x_i), \\ & \geq \limsup_{n \rightarrow \infty} \left[\langle A_i(x_1^n, \dots, y_i, \dots, x_k^n) + S_i(x_1^n, \dots, x_i^n, \dots, x_k^n) - f_i, y_i - x_i^n \rangle_{X_i^* \times X_i} + \right. \\ & \quad \left. J_i^\circ(x_i^n; y_i - x_i^n) + \phi_i(y_i) - \phi_i(x_i^n) \right], \end{aligned}$$

$$\begin{aligned}
&\geq \limsup_{n \rightarrow \infty} \left[\langle A_i(x_1^n, \dots, x_i^n, \dots, x_k^n) + S_i(x_1^n, \dots, x_i^n, \dots, x_k^n) - f_i, y_i - x_i^n \rangle_{X_i^* \times X_i} + \right. \\
&\quad \left. J_i^\circ(x_i^n; y_i - x_i^n) + \phi_i(y_i) - \phi_i(x_i^n) \right], \quad [\text{by } (H_m)] \\
&\geq \limsup_{n \rightarrow \infty} \left[-\rho_n \|y_i - x_i^n\|_{X_i} \right], \quad [\text{by } (3.1)], \\
&= 0, \quad \forall y_i \in X_i.
\end{aligned}$$

Now, due to the convexity of ϕ_i , (H_h) and Lemma 1, we can derive (by invoking the analogous proof outline of Lemma 2) that

$$\begin{aligned}
&\langle A_i(x_1, \dots, x_i, \dots, x_k) + S_i(x_1, \dots, x_i, \dots, x_k) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) \\
&\quad + \phi_i(y_i) - \phi_i(x_i) \geq 0, \quad \text{for each } i \in \{1, \dots, k\},
\end{aligned}$$

which means that $x \in Q$. For the completeness of the proof, let us assume by contradiction that the SMHVI has two distinct solutions, say, $x, z \in Q$. Since $Q \subset \Upsilon(\rho) \forall \rho > 0$, so as $x, z \in \Upsilon(\rho) \forall \rho > 0$. Also, since $\text{diam}(\Upsilon(\rho)) \rightarrow 0$ as $\rho \rightarrow 0$, it is straightforward to deduce that

$$\|x - z\| \leq \text{diam}(\Upsilon(\rho)) \rightarrow 0,$$

which contradicts $x \neq z$, and this confirms the uniqueness of the solution. \square

Theorem 4. For each $i \in \{1, \dots, k\}$, let X_i be a real Banach space with dual space X_i^* and assume that $A_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$ satisfies the hypothesis (H_a) , (H_h) and (H_m) . Further, suppose that for each $i \in \{1, \dots, k\}$, $S_i: X_1 \times \dots \times X_i \times \dots \times X_k \rightarrow X_i^*$, $J_i: X_i \rightarrow \mathbb{R}$ and $\phi_i: X_i \rightarrow \mathbb{R} \cup \{\infty\}$ satisfy the hypothesis (H_c) , (H_j) and (H_ϕ) , respectively. Then SMHVI is strongly well-posedness in generalized sense iff

$$\Upsilon(\rho) \neq \emptyset \quad \forall \rho > 0 \quad \text{and} \quad \mu(\Upsilon(\rho)) \rightarrow 0 \quad \text{as } \rho \rightarrow 0. \quad (4.4)$$

Proof. Assume that SMHVI is strongly well-posed in generalized sense. For any $\rho > 0$, then $Q \neq \emptyset$ and $Q \subset \Upsilon(\rho)$. Moreover, we claim that the set Q is compact. For any sequence, say, $x^n \in X_1 \times \dots \times X_i \times \dots \times X_k$ s.t. $x^n \in Q$ and so $x^n \in \Upsilon(\rho)$ for all $\rho > 0$. For any $\rho > 0$, it follows from $Q \subset \Upsilon(\rho)$ that $\{x^n\}$ is an approximating sequence for SMHVI. Since the SMHVI is strongly well-posed in generalized sense, therefore $\{x^n\}$ has a subsequence that converges to some point of the set Q , which indicates that the set Q is compact and hence $\mu(Q) = 0$. Now, we claim that $\mu(\Upsilon(\rho)) \rightarrow 0$. In fact, for any $\rho > 0$, $Q \subset \Upsilon(\rho)$ implies that

$$\mathcal{H}(\Upsilon(\rho), Q) = \max\{e(\Upsilon(\rho), Q), e(Q, \Upsilon(\rho))\} = e(\Upsilon(\rho), Q). \quad (4.5)$$

Also, from the compactness of Q and (4.5), we obtain

$$\mu(\Upsilon(\rho)) \leq 2\mathcal{H}(\Upsilon(\rho), Q) + \mu(Q) \leq 2\mathcal{H}(\Upsilon(\rho), Q) = 2e(\Upsilon(\rho), Q).$$

Hence, to prove $\mu(\Upsilon(\rho)) \rightarrow 0$ as $\rho \rightarrow 0$, it is enough to show that $e(\Upsilon(\rho), Q) \rightarrow 0$ as $\rho \rightarrow 0$. Arguing by contradiction, we assume that $e(\Upsilon(\rho), Q) \not\rightarrow 0$ as $\rho \rightarrow 0$. Thus, there exists a constant $\varepsilon > 0$ and a sequence $\{\rho_n\} \subset \mathbb{R}_+$ with $\rho_n \rightarrow 0$ as $n \rightarrow \infty$ and $w^n \in \Upsilon(\rho_n)$ satisfying

$$d(w^n, Q) > \varepsilon, \quad \forall n \in \mathbb{N}. \quad (4.6)$$

Since $\{w^n\}$ is an approximating sequence for SMHVI and SMHVI is well-posed in generalized sense, therefore, there exists a subsequence $\{w^{n_i}\}$ of $\{w^n\}$, which converges to some point of Q . So we get

$$0 < \varepsilon < d(w^{n_i}, Q) \rightarrow 0,$$

which contradicts (4.6).

Conversely, assume that (4.4) holds ($\Upsilon(\rho) \neq \emptyset$). By Corollary 1, we have that $\Upsilon(\rho)$ is closed for all $\rho > 0$. Next, we claim that

$$Q = \bigcap_{\rho > 0} \Upsilon(\rho).$$

It is evident that $Q \subset \bigcap_{\rho > 0} \Upsilon(\rho)$. Conversely, let $x = (x_1, \dots, x_i, \dots, x_k) \in \bigcap_{\rho > 0} \Upsilon(\rho)$. So, there exists $\rho_n > 0$ s.t. $\rho_n \rightarrow 0$, which leads to $x = (x_1, \dots, x_i, \dots, x_k) \in \Upsilon(\rho_n)$. Thus for each $i \in \{1, \dots, k\}$, we get

$$\begin{aligned} & \langle A_i(x) + S_i(x) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) + \phi_i(y_i) - \phi_i(x_i) \\ & \geq -\rho_n \|y_i - x_i\|_{X_i}, \quad \forall y_i \in X_i. \end{aligned} \quad (4.7)$$

Letting $n \rightarrow \infty$ in (4.7), we get

$$\langle A_i(x) + S_i(x) - f_i, y_i - x_i \rangle_{X_i^* \times X_i} + J_i^\circ(x_i; y_i - x_i) + \phi_i(y_i) - \phi_i(x_i) \geq 0, \quad \forall y_i \in X_i.$$

For any $z = (z_1, \dots, z_i, \dots, z_k) \in X_1 \times \dots \times X_i \times \dots \times X_k$ and $t \in (0, 1)$, let $y_i = x_i + t(z_i - x_i) \in X_i, i \in \{1, \dots, k\}$. Now using the convexity of ϕ_i and Lemma 1, it is easy to draw that $x \in Q$ and hence $\bigcap_{\rho > 0} \Upsilon(\rho) \subset Q$. This shows that $Q = \bigcap_{\rho > 0} \Upsilon(\rho)$. Since $\mu(\Upsilon(\rho)) \rightarrow 0$ as $\rho \rightarrow 0$. By generalized Cantor theorem (see [12]), it follows that Q is nonempty and compact with $\Upsilon(\rho) \rightarrow Q$ as $\rho \rightarrow 0$ w.r.t. \mathcal{H} , i.e.,

$$\mathcal{H}(\Upsilon(\rho), Q) \rightarrow 0 \quad \text{as } \rho \rightarrow 0. \quad (4.8)$$

As,

$$e(\Upsilon(\rho), Q) = \mathcal{H}(\Upsilon(\rho), Q), \quad \text{for } \rho > 0.$$

Hence, from (4.8), we get

$$e(\Upsilon(\rho), Q) \rightarrow 0 \quad \text{as } \rho \rightarrow 0. \quad (4.9)$$

In order to complete the proof, let $x^n = (x_1^n, \dots, x_i^n, \dots, x_k^n) \in X_1 \times \dots \times X_i \times \dots \times X_k$ be an approximating sequence for SMHVI, thus there exists $\{\rho_n\} \subset \mathbb{R}_+$ with $\rho_n \rightarrow 0$ s.t. for each $i \in \{1, \dots, k\}$, we have

$$\begin{aligned} & \langle A_i(x^n) + S_i(x^n) - f_i, y_i - x_i^n \rangle_{X_i^* \times X_i} + J_i^\circ(x_i^n; y_i - x_i^n) + \phi_i(y_i) - \phi_i(x_i^n) \\ & \geq -\rho_n \|y_i - x_i^n\|_{X_i}, \quad \forall y_i \in X_i, \end{aligned}$$

i.e., $x^n \in \Upsilon(\rho_n)$. It ensures from (4.9) that

$$d(x^n, Q) \leq e(\Upsilon(\rho_n), Q) \rightarrow 0.$$

As the solution set Q is compact, so there exists $w^n = (w_1^n, \dots, w_i^n, \dots, w_k^n) \in Q$ s.t.

$$\|x^n - w^n\| = d(x^n, Q) \rightarrow 0. \quad (4.10)$$

Also, from the compactness of Q , the sequence $\{w^n\}$ has a subsequence $\{w^{n_l}\}$ s.t. $w^{n_l} \rightarrow w'$, (for some $w' \in Q$). Hence, it follows from (4.10) that

$$\|x^{n_l} - w'\| \leq \|x^{n_l} - w^{n_l}\| + \|w^{n_l} - w'\| \rightarrow 0,$$

which estimates $x^{n_l} \rightarrow w'$. This derives the strongly well-posedness of the SMHVI. \square

The next theorem is arranged by setting $X_i = \mathbb{R}$ for each $i \in \{1, \dots, k\}$. Hence $X_1 \times \dots \times X_i \times \dots \times X_k = \mathbb{R}^k$. Here, we present some conditions by which the SMHVI is well-posed in generalized sense.

Theorem 5. *For each $i \in \{1, \dots, k\}$, let $A_i: \mathbb{R}^k \rightarrow \mathbb{R}$ be a continuous mapping satisfying the hypothesis (H_h) and (H_m) . Further, assume that for each $i \in \{1, \dots, k\}$, $S_i: \mathbb{R}^k \rightarrow \mathbb{R}$, $J_i: \mathbb{R} \rightarrow \mathbb{R} \cup \{\infty\}$ and $\phi_i: \mathbb{R} \rightarrow \mathbb{R} \cup \{\infty\}$ satisfy the hypothesis (H_c) , (H_j) and (H_ϕ) , respectively. If there exist $n_0 \in \mathbb{N}$ and $\rho > 0$ s.t. $\Upsilon(\rho)$ is nonempty, bounded and for every approximating sequence $\{x^n\}$ of SMHVI $x^n \in \Upsilon(\rho) \forall n > n_0$. Then SMHVI is strongly well-posedness in generalized sense.*

Proof. Let $x^n = (x_1^n, \dots, x_i^n, \dots, x_k^n) \in \mathbb{R}^k$ be an approximating sequence of the SMHVI. Then there exists $\{\rho_n\} \subset \mathbb{R}_+$ s.t. $\rho_n \rightarrow 0$ as $n \rightarrow \infty$. Thus for each $i \in \{1, \dots, k\}$,

$$\begin{aligned} & \langle A_i(x^n) + S_i(x^n) - f_i, y_i - x_i^n \rangle + J_i^\circ(x_i^n; y_i - x_i^n) + \phi_i(y_i) - \phi_i(x_i^n) \\ & \geq -\rho_n \|y_i - x_i^n\|_{\mathbb{R}}, \quad \forall y_i \in \mathbb{R}, \end{aligned} \quad (4.11)$$

On the other hand, let $\rho > 0$ be s.t. $\Upsilon(\rho)$ is nonempty and bounded. Accordingly, there exists $n_0 \in \mathbb{N}$ s.t. $x^n \in \Upsilon(\rho) \forall n > n_0$. This formulates that $\{x^n\}$ is bounded. Thus, there exists a subsequence $\{x^{n_l}\}$ of $\{x^n\}$ s.t. $x^{n_l} = (x_1^{n_l}, \dots, x_i^{n_l}, \dots, x_k^{n_l}) \rightarrow \hat{x} = (\hat{x}_1, \dots, \hat{x}_i, \dots, \hat{x}_k)$. Further, due to the hypothesis (H_c) , (H_m) , (H_j) and the lower semi-continuity of ϕ_i along with Lemma 1, we have the following estimates

$$\begin{aligned} & \langle A_i(\hat{x}_1, \dots, y_i, \dots, \hat{x}_k) + S_i(\hat{x}_1, \dots, \hat{x}_i, \dots, \hat{x}_k) - f_i, y_i - \hat{x}_i \rangle \\ & + J_i^\circ(\hat{x}_i; y_i - \hat{x}_i) + \phi_i(y_i) - \phi_i(\hat{x}_i), \end{aligned}$$

$$\begin{aligned}
&\geq \limsup_{l \rightarrow \infty} \left[\langle A_i(x_1^{n_l}, \dots, x_i, \dots, x_k^{n_l}) + S_i(x_1^{n_l}, \dots, x_i^{n_l}, \dots, x_k^{n_l}) - f_i, y_i - x_i^{n_l} \rangle \right. \\
&\quad \left. + J_i^\circ(x_i^{n_l}; y_i - x_i^{n_l}) + \phi_i(y_i) - \phi_i(x_i^{n_l}) \right], \\
&\geq \limsup_{l \rightarrow \infty} \left[\langle A_i(x_1^{n_l}, \dots, x_i^{n_l}, \dots, x_k^{n_l}) + S_i(x_1^{n_l}, \dots, x_i^{n_l}, \dots, x_k^{n_l}) - f_i, y_i - x_i^{n_l} \rangle \right. \\
&\quad \left. + J_i^\circ(x_i^{n_l}; y_i - x_i^{n_l}) + \phi_i(y_i) - \phi_i(x_i^{n_l}) \right], \\
&\geq \limsup_{l \rightarrow \infty} [-\rho_n \|y_i - x_i^{n_l}\|_{\mathbb{R}}], \quad [\text{by (4.11)}], \\
&= 0.
\end{aligned}$$

By following the arguments similar to Lemma 2, we can deduce (by hemicontinuity of A_i and the convexity of ϕ_i) that

$$\begin{aligned}
&\langle A_i(\hat{x}_1, \dots, \hat{x}_i, \dots, \hat{x}_k) + S_i(\hat{x}_1, \dots, \hat{x}_i, \dots, \hat{x}_k) - f_i, y_i - \hat{x}_i \rangle \\
&\quad + J_i^\circ(\hat{x}_i; y_i - \hat{x}_i) + \phi_i(y_i) - \phi_i(\hat{x}_i) \geq 0,
\end{aligned}$$

which means that $(\hat{x}_1, \dots, \hat{x}_i, \dots, \hat{x}_k) \in Q$. Thus, the SMHVI is strongly well-posed in generalized sense. \square

Remark 3. Proposition 1 generalizes Lemma 3.2 of [22] and Lemma 3.2 of [23]. Further, Theorem 3 and Theorem 4 generalize Theorem 3.1 of [22] and Theorem 3.2 of [22], respectively. Also, if $k = 1$ with $S_1 = \theta_1$ and $\phi_1 = 0$ then Theorem 3 and Theorem 4 reduce to Theorem 3.1 of [23] and Theorem 3.2 of [23], respectively.

ACKNOWLEDGEMENTS

The authors are grateful to the referees for their insightful comments and suggestions, which have greatly enhanced the clarity and coherence of the article.

REFERENCES

- [1] J. Cen, C. Min, G. J. Tang, and V. T. Nguyen, "Generalized well-posedness results for a class of new mixed variational inequalities." *Optimization.*, vol. 72, no. 2, pp. 411–437, 2023, doi: [10.1080/02331934.2021.1970752](https://doi.org/10.1080/02331934.2021.1970752).
- [2] L. C. Ceng, H. Gupta, and C. F. Wen, "Well-posedness by perturbations of variational-hemivariational inequalities with perturbations." *Filomat.*, vol. 26, no. 5, pp. 881–895, 2012, doi: [10.2298/FIL1205881C](https://doi.org/10.2298/FIL1205881C).
- [3] L. C. Ceng, N. Hadjisavvas, S. Schaible, and J. C. Yao, "Well-posedness for mixed quasivariational-like inequalities," *J. Optim. Theory Appl.*, vol. 139, pp. 109–125, 2008, doi: [10.1007/s10957-008-9428-9](https://doi.org/10.1007/s10957-008-9428-9).

- [4] L. C. Ceng and J. C. Yao, “Well-posedness of generalized mixed variational inequalities, inclusion problems and fixed-point problems.” *Nonlinear Anal.*, vol. 69, no. 12, pp. 4585–4603, 2008, doi: [10.1016/j.na.2007.11.015](https://doi.org/10.1016/j.na.2007.11.015).
- [5] L. C. Ceng, Y. C. Liou, J. C. Yao, and Y. Yao, “Well-posedness for systems of time-dependent hemivariational inequalities in Banach spaces.” *J. Nonlinear Sci. Appl.*, vol. 10, no. 8, pp. 4318–4336, 2017, doi: [10.22436/jnsa.010.08.26](https://doi.org/10.22436/jnsa.010.08.26).
- [6] F. H. Clarke, *Optimization and Nonsmooth Analysis*. Philadelphia: SIAM, 1987.
- [7] N. Costea and C. Varga, “Systems of nonlinear hemivariational inequalities and applications.” *Topol. Methods Nonlinear Anal.*, vol. 41, no. 1, pp. 39–65, 2013.
- [8] Y. P. Fang, N. J. Huang, and J. C. Yao, “Well-posedness of mixed variational inequalities inclusion problems and fixed point problems.” *J. Global Optim.*, vol. 41, no. 1, pp. 117–133, 2008, doi: [10.1007/s10898-007-9169-6](https://doi.org/10.1007/s10898-007-9169-6).
- [9] Y. P. Fang, N. J. Huang, and J. C. Yao, “Well-posedness by perturbations of mixed variational inequalities in Banach spaces.” *European J. Oper. Res.*, vol. 201, no. 3, pp. 682–692, 2010, doi: [10.1016/j.ejor.2009.04.001](https://doi.org/10.1016/j.ejor.2009.04.001).
- [10] D. Goeleven and D. Motreanu, “Well-posed hemivariational inequalities.” *Numer. Funct. Anal. Optim.*, vol. 16, no. 7-8, pp. 909–921, 1995, doi: [10.1080/01630569508816652](https://doi.org/10.1080/01630569508816652).
- [11] X. X. Huang and X. Q. Yang, “Generalized Levitin–Polyak well-posedness in constrained optimization.” *SIAM J. Optim.*, vol. 17, no. 1, pp. 243–258, 2006, doi: [10.1137/040614943](https://doi.org/10.1137/040614943).
- [12] K. Kuratowski, “Sur les espaces complets,” *Fund. Math.*, vol. 15, pp. 301–309, 1930, doi: [10.4064/fm-15-1-301-309](https://doi.org/10.4064/fm-15-1-301-309).
- [13] M. B. Lignola, “Well-posedness and L -well-posedness for quasivariational inequalities,” *J. Optim. Theory Appl.*, vol. 128, no. 1, pp. 119–138, 2006, doi: [10.1007/s10957-005-7561-2](https://doi.org/10.1007/s10957-005-7561-2).
- [14] M. B. Lignola and J. Morgan, “Well-posedness for optimization problems with constraints defined by variational inequalities having a unique solution,” *J. Global Optim.*, vol. 16, no. 1, pp. 57–67, 2000, doi: [10.1023/A:1008370910807](https://doi.org/10.1023/A:1008370910807).
- [15] Z. Liu, S. Zeng, and B. Zeng, “Well-posedness for mixed quasi-variational-hemivariational inequalities,” *Topol. Methods Nonlinear Anal.*, vol. 47, no. 2, pp. 561–578, 2016, doi: [10.12775/TMNA.2016.016](https://doi.org/10.12775/TMNA.2016.016).
- [16] R. Lucchetti, *Convexity and well-posed problems*, ser. CMS Books Math./Ouvrages Math. SMC. New York, NY: Springer, 2006, vol. 22, doi: [10.1007/0-387-31082-7](https://doi.org/10.1007/0-387-31082-7).
- [17] R. Lucchetti and F. Patrone, “A characterization of tyhonov well-posedness for minimum problems with applications to variational inequalities.” *Numer. Funct. Anal. Optim.*, vol. 3, no. 4, pp. 461–476, 1981, doi: [10.1080/01630568108816100](https://doi.org/10.1080/01630568108816100).
- [18] S. Migórski, A. Ochal, and M. Sofonea, “Nonlinear inclusions and hemivariational inequalities. Models and analysis of contact problems,” vol. 26, 2013, doi: [10.1007/978-1-4614-4232-5](https://doi.org/10.1007/978-1-4614-4232-5).
- [19] P. D. Panagiotopoulos, “Nonconvex energy functions. Hemivariational inequalities and stationarity principles.” *Acta Mech.*, vol. 48, no. 3–4, pp. 111–130, 1983, doi: [10.1007/BF01170410](https://doi.org/10.1007/BF01170410).
- [20] A. N. Tykhonov, “On the stability of the functional optimization problem,” *USSR J Comput Math Math Phys.*, vol. 6, no. 4, pp. 28–33, 1966, doi: [10.1016/0041-5553\(66\)90003-6](https://doi.org/10.1016/0041-5553(66)90003-6).
- [21] Y. M. Wang, Y. B. Xiao, X. Wang, and Y. J. Cho, “Equivalence of well-posedness between systems of hemivariational inequalities and inclusion problems.” *J. Nonlinear Sci. Appl.*, vol. 9, no. 3, pp. 1178–1192, 2016, doi: [10.22436/jnsa.009.03.44](https://doi.org/10.22436/jnsa.009.03.44).
- [22] Y. B. Xiao and N. J. Huang, “Well-posedness for a class of variational hemivariational inequalities with perturbations.” *J. Optim. Theory Appl.*, vol. 151, no. 1, pp. 33–51, 2011, doi: [10.1007/s10957-011-9872-9](https://doi.org/10.1007/s10957-011-9872-9).
- [23] Y. B. Xiao, N. J. Huang, and M. M. Wong, “Well-posedness of hemivariational inequalities and inclusion problems.” *Taiwanese J. Math.*, vol. 15, no. 3, pp. 1261–1276, 2011, doi: [10.11650/twjm/1500406298](https://doi.org/10.11650/twjm/1500406298).

- [24] Y. B. Xiao, X. Yang, and N. J. Huang, “Some equivalence results for well-posedness of hemivariational inequalities.” *J. Global Optim.*, vol. 61, no. 4, pp. 789–802, 2015, doi: [10.1007/s10898-014-0198-7](https://doi.org/10.1007/s10898-014-0198-7).

Authors’ addresses

Kartikeswar Mahalik

Department of Mathematics, Indian Institute of Technology Kharagpur, Kharagpur-721302, India

E-mail address: kartikiit1234@gmail.com

Chandal Nahak

(**Corresponding author**) Department of Mathematics, Indian Institute of Technology Kharagpur, Kharagpur-721302, India

E-mail address: cnahak@maths.iitkgp.ac.in